

FAMILY HEALTH CENTER PAIN/OPIOID ASSESSMENT NOTE

1. PAIN LOCATION and ETIOLOGY:

2. ANALGESIC REGIMEN:

3. PAIN SCALE (PAST 24 HOURS; 0=NO PAIN to 10=WORST PAIN POSSIBLE)

NOW:
WORST:
BEST:

4. 6 A's:

(1)ANALGESIA: WHAT NUMBER BEST DESCRIBES YOUR PAIN ON AVERAGE IN THE PAST WEEK? (0=No pain to 10=Worst pain possible)

(2)AFFECT: WHAT NUMBER BEST DESCRIBES HOW, DURING THE PAST WEEK, PAIN HAS INTERFERED WITH YOUR ENJOYMENT IN LIFE? (0=Does not interfere to 10=Completely interferes):

(3)ACTIVITY: WHAT NUMBER BEST DESCRIBES HOW, DURING THE PAST WEEK, PAIN HAS INTERFERED WITH YOUR GENERAL ACTIVITY? (0=Does not interfere to 10=Completely interferes):

TOTAL PEG SCORE(1+2+3): BASELINE: LAST WK: TODAY:

(4)ADJUNCTS: WHAT OTHER METHODS ARE YOU USING TO DECREASE YOUR PAIN?

(5)ADVERS

(6)ABERRANT BEHAVIORS:

a. URINE DRUG TEST RESULTS-- DATE:
RESULTS:
REVIEWED and DISCUSSED WITH PATIENT:

b. PILL COUNT: DATE of Rx:
NUMBER OF PILLS FILLED:
NUMBER OF PILLS REMAINING:

c. OTHER: (see Page 2)

UNIVERSAL PRECAUTIONS:

1. GOALS, BENEFITS and RISKS OF TREATMENT discussed with patient and understood

2. CONTROLLED SUBSTANCE AGREEMENT SIGNED ON:

ASSESSMENT and PLAN: _____

PATIENT STICKER HERE

Potential Aberrant Drug-Related Behavior

This section must be completed by the physician

Please check any of the following items that you discovered during your interactions with the patient. Please note that some of these are directly observable (eg, appears intoxicated), while others may require more active listening and/or probing. Use the "Other" section below to note additional details.

- Purposeful over-sedation
- Negative mood change
- Appears intoxicated
- Increasingly unkempt or impaired
- Involvement in car or other accident
- Requests frequent early renewals
- Increased dose without authorization
- Reports lost or stolen prescriptions
- Attempts to obtain prescriptions from other doctors
- Changes route of administration
- Uses pain medication in response to situational stressor
- Insists on certain medications by name
- Contact with street drug culture
- Abusing alcohol or illicit drugs
- Hoarding (ie, stockpiling) of medication
- Arrested by police
- Victim of abuse
- Other: _____
